

## Four Things You Can Do to Prevent Falls:

### ① **Speak up.**

Talk openly with your healthcare provider about fall risks and prevention. Ask your doctor or pharmacist to review your medicines.

### ② **Keep moving.**

Begin an exercise program to improve your leg strength and balance.

### ③ **Get an annual eye exam.**

Replace eyeglasses as needed.

### ④ **Make your home safer.**

Remove clutter and tripping hazards.

## Learn More

Contact your local community or senior center for information on exercise, fall prevention programs, and options for improving home safety, or visit:

- [go.usa.gov/xN9XA](http://go.usa.gov/xN9XA)
- [www.stopfalls.org](http://www.stopfalls.org)



1 in 4 people 65 and older falls each year.

**For more Information Contact:**  
Dawn Delahunty  
[Hello@DawnDelahunty.com](mailto>Hello@DawnDelahunty.com)  
(925) 420 - 4403  
[www.DawnDelahunty.com](http://www.DawnDelahunty.com)

This brochure was produced in collaboration with the following organizations: VA Greater Los Angeles Healthcare System, Geriatric Research Education & Clinical Center (GRECC), and the Fall Prevention Center of Excellence



**Centers for Disease Control and Prevention**  
National Center for Injury Prevention and Control

# Stay Independent

## Learn more about fall prevention.



**STEADI**

## Check Your Risk for Falling

Circle “Yes” or “No” for each statement below			Why it matters
Yes (2)	No (0)	I have fallen in the past year.	People who have fallen once are likely to fall again.
Yes (2)	No (0)	I use or have been advised to use a cane or walker to get around safely.	People who have been advised to use a cane or walker may already be more likely to fall.
Yes (1)	No (0)	Sometimes I feel unsteady when I am walking.	Unsteadiness or needing support while walking are signs of poor balance.
Yes (1)	No (0)	I steady myself by holding onto furniture when walking at home.	This is also a sign of poor balance.
Yes (1)	No (0)	I am worried about falling.	People who are worried about falling are more likely to fall.
Yes (1)	No (0)	I need to push with my hands to stand up from a chair.	This is a sign of weak leg muscles, a major reason for falling.
Yes (1)	No (0)	I have some trouble stepping up onto a curb.	This is also a sign of weak leg muscles.
Yes (1)	No (0)	I often have to rush to the toilet.	Rushing to the bathroom, especially at night, increases your chance of falling.
Yes (1)	No (0)	I have lost some feeling in my feet.	Numbness in your feet can cause stumbles and lead to falls.
Yes (1)	No (0)	I take medicine that sometimes makes me feel light-headed or more tired than usual.	Side effects from medicines can sometimes increase your chance of falling.
Yes (1)	No (0)	I take medicine to help me sleep or improve my mood.	These medicines can sometimes increase your chance of falling.
Yes (1)	No (0)	I often feel sad or depressed.	Symptoms of depression, such as not feeling well or feeling slowed down, are linked to falls.
<b>Total</b>		Add up the number of points for each “yes” answer. If you scored 4 points or more, you may be at risk for falling.	